

Computations of quandle cocycle invariants of surface-links using marked graph diagrams

Seiichi Kamada, Jieon Kim and Sang Youl Lee

*Department of Mathematics, Osaka City University,
Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan*

E-mail: skamada@sci.osaka-cu.ac.jp,

*Department of Mathematics, Graduate School of Natural Sciences,
Pusan National University,*

Busan 609-735, Republic of Korea

E-mail: jieonkim@pusan.ac.kr

Department of Mathematics, Pusan National University,

Busan 609-735, Republic of Korea

E-mail: sangyoul@pusan.ac.kr

February 6, 2015

Abstract

By using the cohomology theory of quandles, quandle cocycle invariants and shadow quandle cocycle invariants are defined for oriented links and surface-links via broken surface diagrams. By using symmetric quandles, symmetric quandle cocycle invariants are also defined for unoriented links and surface-links via broken surface diagrams. A marked graph diagram is a link diagram possibly with 4-valent vertices equipped with markers. S. J. Lomonaco, Jr. and K. Yoshikawa introduced a method of describing surface-links by using marked graph diagrams. In this paper, we give interpretations of these quandle cocycle invariants in terms of marked graph diagrams, and introduce a method of computing them from marked graph diagrams.

1 Introduction

A *surface-link* is a closed 2-manifold smoothly (or piecewise linearly and locally flatly) embedded in the Euclidian 4-space $\mathbb{R}^4$. Two surface-links $\mathcal{L}$

and $\mathcal{L}'$ are said to be *equivalent* if there exists an orientation preserving homeomorphism $h : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ such that $h(\mathcal{L}) = \mathcal{L}'$. When $\mathcal{L}$ and $\mathcal{L}'$ are oriented, it is assumed that $h|_{\mathcal{L}} : \mathcal{L} \rightarrow \mathcal{L}'$ is also an orientation preserving homeomorphism.

A *broken surface diagram* of a surface-link is a projection image in $\mathbb{R}^3$ with over/under sheet information at each double point curve. It is known that two broken surface diagrams present equivalent surface-links if and only if they are related by a finite sequence of Roseman moves (cf. [29]).

A marked graph diagram is a link diagram possibly with 4-valent vertices equipped with markers. S. J. Lomonaco, Jr. [27] and K. Yoshikawa [32] introduced a method of describing surface-links by using marked graph diagrams. Yoshikawa introduced local moves on marked graph diagrams, which are so-called *Yoshikawa moves*. Two marked graph diagrams present equivalent surface-links if and only if they are related by a finite sequence of Yoshikawa moves ([21, 23, 31]). So one can use marked graph diagrams for studying surface-links and their invariants (cf. [1, 13, 14, 22, 23, 24, 25, 26, 30]).

A *quandle* is a set X with a binary operation $* : X \times X \rightarrow X$ satisfying certain conditions derived from Reidemeister moves for classical link diagrams ([15, 28]). By using the cohomology theory of quandles ([5, 7, 8, 9, 10, 12]), *quandle cocycle invariants* and *shadow quandle cocycle invariants* are defined for oriented links and surface-links via broken surface diagrams ([4, 5, 6]). On the other hand, by using symmetric quandles, *symmetric quandle cocycle invariants* are also defined for unoriented links and surface-links via broken surface diagram ([17, 18]). These invariants for surface-links are defined as state-sums over all quandle colorings of sheets and corresponding Boltzman weights that are evaluations of a cocycle at triple points in broken surface diagrams.

The aim of this paper is to interpret of these quandle cocycle invariants in terms of marked graph diagrams, and introduce a method of computing the quandle cocycle invariants from marked graph diagrams.

This paper is organized as follows: In Section 2, we prepare some preliminaries about broken surface diagrams and marked graph diagrams. Section 3 contains a review of quandle cocycle invariants of oriented surface-links. In Section 4, we describe quandle cocycle invariants via marked graph diagrams and give a method of computing the quandle cocycle invariants from marked graph diagrams. Section 5 contains shadow colorings and shadow quandle cocycle invariants of oriented surface-links. In Section 6, we describe how to compute shadow quandle cocycle invariants from marked graph diagrams. In Section 7, we recall symmetric quandles and symmetric quandle cocy-

cle invariants of unoriented surface-links. Section 8 is devoted to giving a method of computing symmetric quandle cocycle invariants from marked graph diagrams.

2 Marked graph diagrams of surface-links


In this section, we recall broken surface diagrams and marked graph diagrams presenting surface-links.

Let $\mathcal{L}$ be a surface-link. By deforming $\mathcal{L}$ by an ambient isotopy of $\mathbb{R}^4$ if necessary, we may assume that the restriction map $q|_{\mathcal{L}} : \mathcal{L} \rightarrow \mathbb{R}^3$ is a general position map, where $q : \mathbb{R}^4 \rightarrow \mathbb{R}^3$ denotes the projection $(x, y, z, t) \mapsto (x, y, t)$. Along the double point curves, one of the sheets (called the *over-sheet*) lies above the other (*under-sheet*) with respect to the z -coordinate. The under-sheets are coherently broken in the projection. The union $\mathcal{B}$ of such broken surfaces is called a *broken surface diagram* of $\mathcal{L}$. When $\mathcal{L}$ is an oriented surface-link, we assume that the sheets of are co-oriented such that the pair (orientation, co-orientation) matches the given (right-handed) orientation of $\mathbb{R}^3$. In [29], D. Roseman introduced seven moves of broken surface diagrams, called *Roseman moves*. Two surface-links are equivalent if and only if their broken surface diagrams are related by a finite sequence of Roseman moves. For more details, see [2, 29].

A *marked graph* is a spatial graph G in $\mathbb{R}^3$ which satisfies the following:

- (1) G is a finite regular graph with 4-valent vertices, say $v_1, v_2, \dots, v_n$.
- (2) Each v_i is a rigid vertex; that is, we fix a rectangular neighborhood N_i homeomorphic to $\{(x, y) | -1 \leq x, y \leq 1\}$, where v_i corresponds to the origin and the edges incident to v_i are represented by $x^2 = y^2$.
- (3) Each v_i has a *marker*, which is the interval on N_i given by $\{(x, 0) | -1 \leq x \leq 1\}$.

Two marked graphs are said to be *equivalent* if they are ambient isotopic in $\mathbb{R}^3$ with keeping the rectangular neighborhoods and markers.

An *orientation* of a marked graph G is a choice of an orientation for each edge of G such that every vertex in G looks like . A marked graph G is said to be *orientable* if it admits an orientation. Otherwise, it is said to be *non-orientable*. Figure 1 shows an oriented marked graph and a




Figure 1: Marked graphs

non-orientable marked graph. Marked graphs can be described by diagrams on $\mathbb{R}^2$ with some 4-valent vertices equipped with markers.

A surface-link $\mathcal{L}$ in $\mathbb{R}^4 = \mathbb{R}^3 \times \mathbb{R}$ can be described in terms of its *cross-sections* $\mathcal{L}_t = \mathcal{L} \cap \mathbb{R}^3 \times \{t\}$, $t \in \mathbb{R}$ (cf. [11]). Let $p : \mathbb{R}^4 \rightarrow \mathbb{R}$ be the projection given by $p(x_1, x_2, x_3, x_4) = x_4$, and we denote by $p_{\mathcal{L}} : \mathcal{L} \rightarrow \mathbb{R}$ the restriction to $\mathcal{L}$. It is known ([19, 20, 27]) that any surface-link $\mathcal{L}$ is equivalent to a surface-link $\mathcal{L}'$, called a *hyperbolic splitting* of $\mathcal{L}$, such that the projection $p_{\mathcal{L}'} : \mathcal{L}' \rightarrow \mathbb{R}$ satisfies that all critical points are non-degenerate, all the index 0 critical points (minimal points) are in $\mathbb{R}^3 \times \{-1\}$, all the index 1 critical points (saddle points) are in $\mathbb{R}^3 \times \{0\}$, and all the index 2 critical points (maximal points) are in $\mathbb{R}^3 \times \{1\}$.

Let $\mathcal{L}$ be a surface-link and let $\mathcal{L}'$ be a hyperbolic splitting of $\mathcal{L}$. The cross-section $\mathcal{L}'_0 = \mathcal{L}' \cap \mathbb{R}^3 \times \{0\}$ at $t = 0$ is a 4-valent graph in $\mathbb{R}^3 \times \{0\}$. We give a marker at each 4-valent vertex (saddle point) that indicates how the saddle point opens up above as illustrated in Figure 2. The resulting marked graph G is called a *marked graph* presenting $\mathcal{L}$. As usual, G is described by a diagram Γ on $\mathbb{R}^2$ which is a generic projection on $\mathbb{R}^2$ with over/under crossing information for each double point such that the restriction to a rectangular neighborhood of each marked vertex is an embedding. Such a diagram is called a *marked graph diagram* or a *ch-diagram* (cf. [30]) presenting $\mathcal{L}$.

When $\mathcal{L}$ is an oriented surface-link, we assume that $\mathcal{L}'_0$ has the induced orientation as the boundary of the oriented surface $\mathcal{L}' \cap (\mathbb{R}^3 \times (-\infty, 0])$.

Let Γ be a marked graph diagram and Γ_0 the singular link diagram obtained from Γ by removing all markers. Let $V(\Gamma) = \{v_1, v_2, \dots, v_n\}$ be the set of all vertices of Γ . For each i ($i = 1, \dots, n$), consider four points v_i^1, v_i^2, v_i^3 , and v_i^4 on Γ in a neighborhood of v_i as in Figure 3. We define

$$\Gamma_+ = [\Gamma_0 \setminus \bigcup_{i=1}^n (\bigcup_{j=1}^4 |v_i, v_i^j|)] \cup [\bigcup_{i=1}^n (|v_i^1, v_i^2| \cup |v_i^3, v_i^4|)],$$




Figure 2: Marking of a vertex

which is called the *positive resolution* of Γ , and

$$\Gamma_- = [\Gamma_0 \setminus \bigcup_{i=1}^n (\bigcup_{j=1}^4 |v_i, v_i^j|)] \cup [\bigcup_{i=1}^n (|v_i^1, v_i^3| \cup |v_i^2, v_i^4|)],$$

the *negative resolution* of Γ , where $|v, w|$ is the line segment connecting v and w . When both resolutions Γ_- and Γ_+ are diagrams of trivial links, we call Γ *admissible*.




Figure 3: v_i^1, v_i^2, v_i^3 , and v_i^4

When Γ is admissible, we construct a surface-link as follows (cf. [17, 19, 20, 32]). Let L_0 denote a graph in $\mathbb{R}^3$ whose diagram is Γ_0 . Let w_i^j and w_i be points on L_0 such that $\pi(w_i^j) = v_i^j$, $\pi(w_i) = v_i$, respectively, where $\pi : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ is the projection $(x, y, z) \mapsto (x, y)$. For each $t \in [0, 1]$, let $w_i^j(t)$ be the point $(1 - t)w_i + tw_i^j \in \mathbb{R}^3$.

For each $t \in [0, 1]$, let L_t^+ be a link defined by

$$L_t^+ = [L_0 \setminus \bigcup_{i=1}^n (\bigcup_{j=1}^4 |w_i, w_i^j(t)|)] \cup [\bigcup_{i=1}^n (|w_i^1(t), w_i^2(t)| \cup |w_i^3(t), w_i^4(t)|)],$$

and for each $t \in [-1, 0]$, let L_t^- be a link defined by

$$L_t^- = [L_0 \setminus \bigcup_{i=1}^n (\bigcup_{j=1}^4 |w_i, w_i^j(-t)|)] \cup [\bigcup_{i=1}^n (|w_i^1(-t), w_i^3(-t)| \cup |w_i^2(-t), w_i^4(-t)|)].$$

Put $L_+ = L_1^+$ and $L_- = L_{-1}^-$. Then L_+ and L_- have diagrams Γ_+ and Γ_- , respectively. Let $B_1^+, \dots, B_\mu^+$ be mutually disjoint 2-disks in $\mathbb{R}^3$ with $\partial(B_1^+ \cup \dots \cup B_\mu^+) = L_+$, and let $B_1^-, \dots, B_\lambda^-$ be mutually disjoint 2-disks in $\mathbb{R}^3$ with $\partial(B_1^- \cup \dots \cup B_\lambda^-) = L_-$.

Let $F(\Gamma)$ be a surface-link in $\mathbb{R}^4 = \mathbb{R}^3 \times \mathbb{R}$ defined by

$$\begin{aligned} F(\Gamma) = & (B_1^- \cup \dots \cup B_\lambda^-) \times \{-2\} \cup L_- \times (-2, -1) \\ & \cup (\cup_{t \in [-1, 0)} L_t^- \times \{t\}) \cup L_0 \times \{0\} \cup (\cup_{t \in (0, 1]} L_t^+ \times \{t\}) \\ & \cup L_+ \times (1, 2) \cup (B_1^+ \cup \dots \cup B_\mu^+) \times \{2\}. \end{aligned}$$

We say that $F(\Gamma)$ is a *surface-link associated to Γ* . It is uniquely determined from Γ up to equivalence (see [19]).

A surface-link $\mathcal{L}$ is said to be *presented* by a marked graph diagram Γ if $\mathcal{L}$ is equivalent to the surface-link $F(\Gamma)$. Any surface-link can be presented by an admissible marked graph diagram. Two admissible marked graph diagrams present equivalent surface-links if and only if they are related by a finite sequence of Yoshikawa moves ([21, 23, 31]).

S. Ashihara introduced a method of constructing a broken surface diagram of a surface-link from its marked graph diagram [1]. For our later use, we describe here his construction. In what follows, by $D \rightarrow D'$ we mean that a link diagram D' is obtained from a link diagram D by a single Reidemeister move (Figure 4) or an ambient isotopy of $\mathbb{R}^2$.



Figure 4: Reidemeister moves of type R_1 , R_2 and R_3

Let Γ be an admissible marked graph diagram, and let Γ_+ and Γ_- be the positive and the negative resolutions.

Since Γ_+ is a diagram of a trivial link, there is a sequence of link diagrams from Γ_+ to a trivial link diagram O related by ambient isotopies of $\mathbb{R}^2$ and Reidemeister moves:

$$\Gamma_+ = D_1 \rightarrow D_2 \rightarrow \dots \rightarrow D_r = O.$$

For each i ($i = 1, \dots, r-1$), let $\{f_t^{(i)}\}_{t \in I}$ be a 1-parameter family of homeomorphisms from $\mathbb{R}^3$ to $\mathbb{R}^3$ which satisfies

$$f_0^{(i)} = \text{id}, \quad f_1^{(i)}(L(D_i)) = L(D_{i+1}),$$

where $L(D_i)$ denotes a link in $\mathbb{R}^3$ with diagram D_i for i ($i = 1, \dots, r$). Without loss of generality, we may assume that $L(D_1) = L_+$ and the following two conditions are satisfied.

- When the move $D_i \rightarrow D_{i+1}$ is an ambient isotopy of $\mathbb{R}^2$, let $\{h_t^{(i)}\}_{t \in I}$ be an ambient isotopy of $\mathbb{R}^2$ such that $h_1^{(i)}(D_i) = D_{i+1}$. Then $f_t^{(i)}$ satisfies $\pi(f_t^{(i)}(L(D_i))) = h_t^{(i)}(\pi(L(D_i)))$ for $t \in I$.
- When the move $D_i \rightarrow D_{i+1}$ is a Reidemeister move, let $B_{(i)}$ be a disk in $\mathbb{R}^2$ where the move is applied, and let $M_{(i)}$ be the subset of $B_{(i)} \times I \subset \mathbb{R}^3$ determined by $\pi(M_{(i)} \cap (B_{(i)} \times \{t\})) = \pi(f_t^{(i)}(L(D_i))) \cap B_{(i)}$ for $t \in I$. Then $M_{(i)}$ is as in Figure 5, 6, or 7.




Figure 5: $M_{(i)}$ for R_1




Figure 6: $M_{(i)}$ for R_2




Figure 7: $M_{(i)}$ for R_3

Take real numbers $t_1, \dots, t_r$ with $1 < t_1 < \dots < t_r < 2$. For each i ($i = 1, \dots, r-1$), we define a homeomorphism $F^{(i)} : \mathbb{R}^4 (= \mathbb{R}^3 \times \mathbb{R}) \rightarrow \mathbb{R}^4$ by

$$F^{(i)}(x, t) = \begin{cases} (x, t) & (t \leq t_i), \\ (f_{\phi(t)}^{(i)}(x), t) & (t_i < t < t_{i+1}), \\ (f_1^{(i)}(x), t) & (t \geq t_{i+1}), \end{cases}$$

where $\phi(t) = (t - t_i)/(t_{i+1} - t_i)$.

Similarly, consider a sequence of link diagrams from Γ_- to a trivial link diagram O' , related by ambient isotopies of $\mathbb{R}^2$ and Reidemeister moves:

$$\Gamma_- = D'_1 \rightarrow D'_2 \rightarrow \dots \rightarrow D'_s = O'.$$

For each j ($j = 1, \dots, s-1$), let $\{g_t^{(j)}\}_{t \in I}$ be a 1-parameter family of homeomorphisms from $\mathbb{R}^3$ to $\mathbb{R}^3$ which satisfies

$$g_0^{(j)} = \text{id}, \quad g_1^{(j)}(L(D'_j)) = L(D'_{j+1}).$$

Without loss of generality, we may assume that $L(D'_1) = L_-$ and the following two conditions are satisfied.

- When the move $D'_j \rightarrow D'_{j+1}$ is an ambient isotopy of $\mathbb{R}^2$, let $\{h'_t\}_{t \in I}$ be an ambient isotopy of $\mathbb{R}^2$ such that $h'_1(D'_j) = D'_{j+1}$. Then $g_t^{(j)}$ satisfies $\pi(g_t^{(j)}(L(D'_j))) = h'_t(\pi(L(D'_j)))$ for $t \in I$.
- When the move $D'_j \rightarrow D'_{j+1}$ is a Reidemeister move, let $B'_{(j)}$ be a disk in $\mathbb{R}^2$ where the move is applied, and let $M'_{(j)}$ be the subset of $B'_{(j)} \times I \subset \mathbb{R}^3$ determined by $\pi(M'_{(j)} \cap (B'_{(j)} \times \{t\})) = \pi(g_t^{(j)}(L(D'_j))) \cap B'_{(j)}$ for $t \in I$. Then $M'_{(j)}$ is as in Figure 5, 6, or 7.

Take real numbers $t'_1, \dots, t'_s$ with $-1 > t'_1 > \dots > t'_s > -2$. For each j ($j = 1, \dots, s-1$), we define a homeomorphism $G^{(j)} : \mathbb{R}^4 \rightarrow \mathbb{R}^4$ by

$$G^{(j)}(x, t) = \begin{cases} (x, t) & (t \geq t'_j), \\ (g_{\psi(t)}^{(j)}(x), t) & (t'_{j+1} < t < t'_j), \\ (g_1^{(j)}(x), t) & (t \leq t'_{j+1}), \end{cases}$$

where $\psi(t) = (t'_j - t)/(t'_j - t'_{j+1})$.

Let $F' = G^{(s-1)} \circ G^{(s-2)} \circ \dots \circ G^{(1)} \circ F^{(r-1)} \circ F^{(r-2)} \circ \dots \circ F^{(1)}(F(\Gamma))$. Then F' is equivalent to $F(\Gamma)$.

Let $B_1, \dots, B_\mu$ be mutually disjoint 2-disks in $\mathbb{R}^3$ such that $\partial(B_1 \cup \dots \cup B_\mu) = L(O)$ and $\pi|_{B_1 \cup \dots \cup B_\mu}$ is an embedding. Let $B'_1, \dots, B'_\lambda$ be mutually disjoint 2-disks in $\mathbb{R}^3$ such that $\partial(B'_1 \cup \dots \cup B'_\lambda) = L(O')$ and $\pi|_{B'_1 \cup \dots \cup B'_\lambda}$ is an embedding. Finally we define F to be the surface constructed as follows:

$$F = (B'_1 \cup \dots \cup B'_\lambda) \times \{-2\} \cup (F' \cap (\mathbb{R}^3 \times (-2, 2))) \cup (B_1 \cup \dots \cup B_\mu) \times \{2\}.$$

It is in general position with respect to the projection $q : \mathbb{R}^4 \rightarrow \mathbb{R}^3, (x, y, z, t) \mapsto (x, y, t)$. The broken surface diagram of F obtained from $q(F)$ is called a *broken surface diagram associated to Γ* , and denoted by $\mathcal{B}(\Gamma)$.

3 Quandle cocycle invariants of oriented surface-links

We recall quandle cocycle invariants of oriented surface-links from [5].

A *quandle* is a set X with a binary operation $*$: $X \times X \rightarrow X$ satisfying that (i) for any $x \in X$, $x * x = x$, (ii) for any $x, y \in X$, there is a unique $u \in X$ such that $x = u * y$, and (iii) for any $x, y, z \in X$, $(x * y) * z = (x * z) * (y * z)$. In (ii), the unique element u is denoted by $x * \bar{y}$, and then $x = u * y = (x * \bar{y}) * y$.

Example 3.1. (1) The *dihedral quandle* of order n is the set $R_n = \{0, 1, \dots, n-1\}$ with the binary operation $i * j = 2j - i \pmod{n}$ for each $i, j \in R_n$.

(2) Let $S_4 = \{0, 1, 2, 3\}$. Define a binary operation $*$: $S_4 \times S_4 \rightarrow S_4$ by

$*$	0	1	2	3
0	0	2	3	1
1	3	1	0	2
2	1	3	2	0
3	2	0	1	3

Then S_4 is a quandle, which is called the *tetrahedral quandle*.

(3) Let G be a group. The *conjugation quandle*, denoted by $\text{conj}(G)$, is G with the operation $x * y = y^{-1}xy$.

Let X be a quandle. For each positive integer n , let $C_n^R(X)$ be the free abelian group generated by n -tuples $(x_1, \dots, x_n)$ of elements of X . We assume $C_n^R(X) = \{0\}$ for $n \leq 0$. Define a homomorphism $\partial_n : C_n^R(X) \rightarrow C_{n-1}^R(X)$ by

$$\begin{aligned} \partial_n(x_1, x_2, \dots, x_n) = & \sum_{i=2}^n (-1)^i [(x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \\ & - (x_1 * x_i, x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)] \end{aligned}$$

for $n \geq 2$ and $\partial_n = 0$ for $n \leq 1$. Then $C_*^R(X) = \{C_n^R(X), \partial_n\}$ is a chain complex. Let $C_n^D(X)$ be the subset of $C_n^R(X)$ generated by n -tuples $(x_1, \dots, x_n)$ with $x_i = x_{i+1}$ for some $i \in 1, \dots, n-1$ if $n \geq 2$; otherwise let $C_n^D(X) = 0$. Then $C_*^D(X) = \{C_n^D(X), \partial_n\}$ is a sub-complex of $C_*^R(X)$. Consider the quotient chain complex $C_*^Q(X) = \{C_n^Q(X), \partial_n\}$, where $C_n^Q(X) = C_n^R(X)/C_n^D(X)$. For an abelian group A , we define chain and cochain complexes by $C_*^Q(X; A) = C_*^Q(X) \otimes A$ and $C_Q^*(X; A) = \text{Hom}(C_*^Q(X), A)$. The homology and cohomology groups are denoted by $H_n^Q(X; A)$ and $H_Q^n(X; A)$, respectively. The cycle and boundary groups (or cocycle and coboundary groups, resp.) are denoted by $Z_n^Q(X; A)$ and $B_n^Q(X; A)$ (or $Z_Q^n(X; A)$ and $B_Q^n(X; A)$, resp.). We will omit the coefficient group A as usual if $A = \mathbb{Z}$.

A homomorphism $\theta : C_3^R(X) \rightarrow A$ is regarded as a 3-cocycle of the cochain complex $C_Q^*(X; A)$, called a *quandle 3-cocycle*, if and only if θ satisfies the following two conditions (where A is written multiplicative):

- $\theta(x, x, y) = 1$ and $\theta(x, y, y) = 1$ for all $x, y \in X$, where 1 is the identity element in A .
- $\theta(x, z, w)\theta(x, y, z)\theta(x * z, y * z, w) = \theta(x, y, w)\theta(x * y, z, w)\theta(x * w, y * w, z * w)$ for each $x, y, z, w \in X$.

Let $\mathcal{B}$ be a broken surface diagram of an oriented surface-link $\mathcal{L}$, and let $S(\mathcal{B})$ be the set of sheets of $\mathcal{B}$. Let X be a quandle. A *coloring* of $\mathcal{B}$ by X is a map $\mathcal{C} : S(\mathcal{B}) \rightarrow X$ satisfying the condition that at each double point curve, if the co-orientation of the over-sheet y is from the under-sheet x to z , then $\mathcal{C}(z) = \mathcal{C}(x) * \mathcal{C}(y)$. See the left of Figure 8. Let $\text{Col}_X(\mathcal{B})$ denote the set of all colorings of $\mathcal{B}$ by X .




Figure 8: A double point curve and a triple point

Let τ be a triple point of $\mathcal{B}$. The *sign* of τ is positive if the co-orientations

of the top, the middle and the bottom sheets at τ in this order match the given (right-handed) orientation of $\mathbb{R}^3$. Otherwise, the sign is negative. There are eight complementary regions of $\mathcal{B}$ around τ . (Some of them may be the same.) There is a unique region such that the co-orientations of the sheets facing the region point from the region to the opposite regions. We call this region the *source region* of τ .

For a 3-cocycle $\theta \in Z_Q^3(X; A)$, the quandle cocycle invariant $\Phi_\theta(\mathcal{L})$ of an oriented surface-link $\mathcal{L}$ associated to θ is defined as follows. Let $\mathcal{B}$ be a broken surface diagram of $\mathcal{L}$. Let $\mathcal{C} : S(\mathcal{B}) \rightarrow X$ be a coloring of $\mathcal{B}$. Let τ be a triple point of $\mathcal{B}$ and let x_1, x_2 , and x_3 be colors of the bottom, the middle, and the top sheets facing the source region of τ , respectively. Let $\epsilon(\tau)$ denote the sign of τ . See Figure 8, where $\epsilon(\tau) = 1$. The (*Boltzman*) *weight* $B_\theta(\tau, \mathcal{C})$ at τ with respect to $\mathcal{C}$ is defined to be

$$B_\theta(\tau, \mathcal{C}) = \theta(x_1, x_2, x_3)^{\epsilon(\tau)}.$$

The *partition function* or *state-sum* of $\mathcal{B}$ (associated to θ) is

$$\Phi_\theta(\mathcal{B}) = \sum_{\mathcal{C} \in \text{Col}_X(\mathcal{B})} \prod_{\tau \in T(\mathcal{B})} B_\theta(\tau, \mathcal{C}) \in \mathbb{Z}[A],$$

where $T(\mathcal{B})$ is the set of all triple points in $\mathcal{B}$.

Theorem 3.2 ([5]). *Let $\mathcal{L}$ be an oriented surface-link and let $\mathcal{B}$ be a broken surface diagram of $\mathcal{L}$. The partition function $\Phi_\theta(\mathcal{B})$ does not depend on the choice of $\mathcal{B}$. Thus it is an invariant of $\mathcal{L}$.*

We call $\Phi_\theta(\mathcal{B})$ the *quandle cocycle invariant* of $\mathcal{L}$ associated to θ , and denote it by $\Phi_\theta(\mathcal{L})$.

4 How to compute quandle cocycle invariants from marked graph diagrams

In this section we introduce a method of computing quandle cocycle invariants from marked graph diagrams.

Let Γ be an oriented marked graph diagram and let $V(\Gamma)$ denote the set of all marked vertices of Γ . By an *arc* of Γ we mean a connected component of $\Gamma \setminus V(\Gamma)$. (At a crossing of Γ the under-arcs are assumed to be cut.) Let $A(\Gamma)$ denote the set of arcs of Γ . Since Γ is oriented, we assume that it is co-oriented: The co-orientation of an arc of Γ satisfies that the pair (orientation, co-orientation) matches the (right-handed) orientation of the

plane. At a crossing, if the pair of the co-orientation of the over-arc and that of the under-arc matches the (right-handed) orientation of the plane, then the crossing is called *positive*; otherwise it is *negative*. The crossing in (a) of Figure 9 is positive and that in (b) is negative.

Definition 4.1. Let X be a quandle and let Γ be an oriented marked graph diagram. A *coloring* of Γ by X is a map $\mathcal{C} : A(\Gamma) \rightarrow X$ satisfying the following conditions (1) and (2):

- (1) For each crossing c , let s_2 be the over-arc and let s_1 and s_3 be the under-arcs as shown in (a) or (b) of Figure 9 such that the co-orientation of s_2 points from s_1 to s_3 . Then $\mathcal{C}(s_3) = \mathcal{C}(s_1) * \mathcal{C}(s_2)$.

(In this case, s_1 is called the *source arc* and s_3 is called the *target arc* at c . The quandle element $\mathcal{C}(s_i)$ is called a *color* of the arc s_i .)

- (2) For each marked vertex v , let s_1, s_2, s_3 and s_4 be the arcs of Γ as shown in (c) or (d) of Figure 9. Then $\mathcal{C}(s_1) = \mathcal{C}(s_2) = \mathcal{C}(s_3) = \mathcal{C}(s_4)$.




Figure 9: Labels at a crossing

We denote by $\text{Col}_X(\Gamma)$ the set of colorings of Γ by X .

Theorem 4.2. Let $\mathcal{L}$ be an oriented surface-link. Let Γ and $\mathcal{B}$ be a marked graph diagram and a broken surface diagram presenting $\mathcal{L}$, respectively. Then there is a bijection from $\text{Col}_X(\Gamma)$ to $\text{Col}_X(\mathcal{B})$.

Proof. The fundamental quandle $Q(\Gamma)$ is defined by a quandle generated by $A(\Gamma)$ and the defining relations $s_3 = s_1 * s_2$ for s_1, s_2, s_3 as in (a) or (b) in Figure 9 and $s_1 = s_2 = s_3 = s_4$ for $s_1, \dots, s_4$ as in (c) or (d). Without loss of generality, we may assume that $\mathcal{B}$ is a broken surface diagram associated to Γ . Then by the same argument with [1] we see that there is a natural isomorphism from $Q(\Gamma)$ to the fundamental quandle $Q(\mathcal{B})$ of $\mathcal{B}$. Since $\text{Col}_X(\Gamma)$ is identified with $\text{Hom}(Q(\Gamma), X)$ and $\text{Col}_X(\mathcal{B})$ is identified with $\text{Hom}(Q(\mathcal{B}), X)$, we have a bijection from $\text{Col}_X(\Gamma)$ to $\text{Col}_X(\mathcal{B})$. $\square$

Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$ and Γ_+ the positive resolution of Γ . Let $\Gamma_+ = D_1 \rightarrow D_2 \rightarrow \cdots \rightarrow D_r = O$ be a sequence of link diagrams from Γ_+ to a trivial link diagram O related by ambient isotopies of $\mathbb{R}^2$ and oriented Reidemeister moves. Let $I_+^3 = \{i \mid D_i \rightarrow D_{i+1} \text{ is a move of type } R_3\}$. For each $i \in I_+^3$, let $B_{(i)}$ be a disk in $\mathbb{R}^2$ where the move $D_i \rightarrow D_{i+1}$ is applied.

Similarly, let Γ_- be the negative resolution of Γ and $\Gamma_- = D'_1 \rightarrow D'_2 \rightarrow \cdots \rightarrow D'_s = O'$ a sequence of link diagrams from Γ_- to a trivial link diagram O' related by ambient isotopies of $\mathbb{R}^2$ and Reidemeister moves. Let $I_-^3 = \{j \mid D'_j \rightarrow D'_{j+1} \text{ is a move of type } R_3\}$. For each $j \in I_-^3$, let $B'_{(j)}$ be a disk in $\mathbb{R}^2$ where the move $D'_j \rightarrow D'_{j+1}$ is applied.

We define two functions ϵ_{tm} and ϵ_b from the disjoint union $I_+^3 \amalg I_-^3$ to $\{\pm 1\}$ as follows:

Let $i \in I_+^3$ (or $i \in I_-^3$, resp.) and let c be the crossing between the top arc and the two middle arcs in $D_i \cap B_{(i)}$ (or $D'_i \cap B'_{(i)}$, resp.) and let n_1 be the co-orientation of the bottom arc. Define $\epsilon_{tm}(i)$ and $\epsilon_b(i)$ for $i \in I_+^3 \amalg I_-^3$ by

$$\epsilon_{tm}(i) = \text{sign}(c), \quad (4.1)$$

$$\epsilon_b(i) = \begin{cases} 1 & \text{if } n_1 \text{ points from } c, \\ -1 & \text{otherwise.} \end{cases} \quad (4.2)$$

Definition 4.3. Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$. Let $\mathcal{C} : A(\Gamma) \rightarrow X$ be a coloring of Γ and let $\theta \in Z_Q^3(X; A)$.

Let $i \in I_+^3 \amalg I_-^3$. Let R be the source region of the crossing c , i.e., the quadrant from which all co-orientations of the top arc and the middle arc point outwards. Let R' be the opposite region of R with respect to the top arc. The (*Boltzman*) *weight* $B_\theta(i, \mathcal{C})$ at i with respect to $\mathcal{C}$ is defined by

$$B_\theta(i, \mathcal{C}) = \theta(x_1, x_2, x_3)^{\epsilon_{tm}(i)\epsilon_b(i)},$$

where x_2 and x_3 are the colors of the middle arc and the top semi-arc facing R , respectively, and x_1 is the color of the bottom semi-arc which is in R or x_1 is the element with $x_1 = a * \overline{x_3}$, where a is the color of the bottom semi-arc which is in R' . See Figure 10.

Definition 4.4. Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$. The *partition function* or *state-sum* of Γ (associated to θ) is

$$\Phi_\theta(\Gamma) = \sum_{\mathcal{C} \in \text{Col}_X(\Gamma)} \left(\prod_{i \in I_+^3} B_\theta(i, \mathcal{C}) \prod_{j \in I_-^3} B_\theta(j, \mathcal{C})^{-1} \right).$$




Figure 10: A (Boltzman) weight at $i \in I_+^3 \amalg I_-^3$

Theorem 4.5. *Let $\mathcal{L}$ be an oriented surface-link and Γ a marked graph diagram of $\mathcal{L}$. Then for any $\theta \in Z_Q^3(X; A)$, $\Phi_\theta(\mathcal{L}) = \Phi_\theta(\Gamma)$.*

Proof. Let $\mathcal{B} = \mathcal{B}(\Gamma)$ be a broken surface diagram associated to Γ . It is sufficient to prove that $\Phi_\theta(\Gamma) = \Phi_\theta(\mathcal{B})$.

Since there is a natural bijection between $\text{Col}_X(\Gamma)$ and $\text{Col}_X(\mathcal{B})$ (as in the proof of Theorem 4.2), it suffices to show the following claim.

Claim: For each coloring $\mathcal{C} \in \text{Col}_X(\Gamma)$,

$$\prod_{i \in I_+^3} B_\theta(i, \mathcal{C}) \prod_{j \in I_-^3} B_\theta(j, \mathcal{C})^{-1} = \prod_{\tau \in T(\mathcal{B})} B_\theta(\tau, \mathcal{C}),$$

where $\mathcal{C} \in \text{Col}_X(\mathcal{B})$ in the right hand side is the corresponding coloring.

Proof of Claim. Let $\mathcal{B}_j^i = \mathcal{B} \cap (\mathbb{R}^2 \times [t_j', t_i])$ for $i = 1, \dots, r$ and $j = 1, \dots, s$. Let $\phi : (\mathbb{R}^2, \Gamma_0) \rightarrow (\mathbb{R}^2 \times [t_1', t_1], \mathcal{B}_1^1)$ be the natural embedding at $t = 0$ as in Figure 11. The vertices of Γ_0 correspond to the saddle points in $\mathcal{B}_1^1$ and the crossings of Γ_0 correspond to the intersection of $\mathbb{R}^2 \times \{0\}$ and the double point curves in $\mathcal{B}_1^1$. There are no triple points in $\mathcal{B}_1^1$.




Figure 11: $\phi : (\mathbb{R}^2, \Gamma_0) \rightarrow (\mathbb{R}^2 \times [t_1', t_1], \mathcal{B}_1^1)$

Let $\mathcal{B}_i = \mathcal{B} \cap (\mathbb{R}^2 \times [t_i, t_{i+1}])$ for $i = 1, \dots, r-1$ and $\mathcal{B}'_j = \mathcal{B} \cap (\mathbb{R}^2 \times [t'_{j+1}, t'_j])$ for $j = 1, \dots, s-1$. Note that $T(\mathcal{B}) = \left(\bigcup_{i=1}^{r-1} T(\mathcal{B}_i) \right) \cup \left(\bigcup_{j=1}^{s-1} T(\mathcal{B}'_j) \right)$.

If the move $D_i \rightarrow D_{i+1}$ is an ambient isotopy of $\mathbb{R}^2$, then $D_i \times [t_i, t_{i+1}] \cong \mathcal{B}_i$, and there are no triple points in $\mathcal{B}_i$.

Suppose that the move $D_i \rightarrow D_{i+1}$ is a Reidemeister move. Since $D_i \setminus B_{(i)}$ and $D_{i+1} \setminus B_{(i)}$ are identical, there are no triple points in $\mathcal{B}_i \setminus M_{(i)}$ and we have $T(\mathcal{B}_i) = T(M_{(i)})$, where $M_{(i)}$ is a subset of $B_{(i)} \times I$ determined by $\pi(M_{(i)} \cap (B_{(i)} \times \{t\})) = \pi(f_t^{(i)}(L(D_i))) \cap B_{(i)}$ for $t \in I$ and a homeomorphism $f_t^{(i)} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ satisfying $f_0^{(i)} = \text{id}$ and $f_1^{(i)}(L(D_i)) = L(D_{i+1})$.

If the move $D_i \rightarrow D_{i+1}$ is of type R_1 or R_2 , then there are no triple points in $M_{(i)}$. See Figures 5 and 6.

If the move $D_i \rightarrow D_{i+1}$ is of type R_3 , then there is a triple point τ_i in $M_{(i)}$ as in Figure 12 and $T(\mathcal{B}_i) = \{\tau_i\}$. Then $\bigcup_{i=1}^{r-1} T(\mathcal{B}_i) = \{\tau_i \mid i \in I_+^3\}$.




Figure 12: A (Boltzman) weight

Similarly, suppose that the move $D'_j \rightarrow D'_{j+1}$ is a Reidemeister move and $M'_{(j)}$ is a subset of $B'_{(j)} \times I$ determined by $\pi(M'_{(j)} \cap (B'_{(j)} \times \{t\})) = \pi(g_t^{(j)}(L(D'_j))) \cap B'_{(j)}$, where $g_t^{(j)} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is a homeomorphism satisfying $g_0^{(j)} = \text{id}$ and $g_1^{(j)}(L(D'_j)) = L(D'_{j+1})$ for $t \in I$. There is a triple point $\tau'_j \in M'_{(j)}$ for $j \in I_-^3$. We have that $\bigcup_{j=1}^{s-1} T(\mathcal{B}'_j) = \{\tau'_j \mid j \in I_-^3\}$.

Now we have

$$T(\mathcal{B}) = \{\tau_i \mid i \in I_+^3\} \cup \{\tau'_j \mid j \in I_-^3\}. \quad (4.3)$$

Let $i \in I_+^3$, i.e., $D_i \rightarrow D_{i+1}$ is a move of type R_3 and let τ_i be the triple point in $M_{(i)}$. Let n_1 , n_2 and n_3 be the co-orientations of the bottom, the middle and the top arcs of D_i in $B_{(i)}$, respectively. By an ambient isotopy, we deform $M_{(i)}$ in $B_{(i)} \times I$ to the standard form of the neighborhood of the triple point τ_i as in Figure 12. Let $\bar{n}_1$, $\bar{n}_2$, and $\bar{n}_3$ be the normal vectors corresponding to n_1 , n_2 , and n_3 , respectively. Without loss of generality, we may assume $\bar{n}_3 = \mathbf{e}_1$, $\bar{n}_2 = \epsilon \mathbf{e}_2$ and $\bar{n}_1 = \epsilon' \mathbf{e}_3$ for some $\epsilon, \epsilon' \in \{1, -1\}$. Here $\mathbf{e}_1 = (1, 0, 0)$, $\mathbf{e}_2 = (0, 1, 0)$ and $\mathbf{e}_3 = (0, 0, 1)$. See Figure 12.

Let c be the crossing between the top and the middle arcs in $B_{(i)}$. It is clear from Figure 12 that $\epsilon = \text{sign}(c)$. By (4.1), $\epsilon = \text{sign}(c) = \epsilon_{tm}(i)$. Hence $\bar{n}_2 = \epsilon_{tm}(i) \mathbf{e}_2$.

The sign ϵ' depends on the co-orientation n_1 of the bottom arc. If n_1 points from c , then $\epsilon' = 1$. If n_1 points toward c , then $\epsilon' = -1$. So, by (4.2), $\epsilon' = \epsilon_b(i)$ and hence $\bar{n}_1 = \epsilon_b(i) \mathbf{e}_3$.

On the other hand, by definition, the sign $\epsilon(\tau_i)$ of the triple point τ_i is positive if the co-orientations of the top, the middle and the bottom sheets in this order match the given (right-handed) orientation of $\mathbb{R}^3$. Otherwise, the sign $\epsilon(\tau_i)$ is negative. This gives

$$\epsilon(\tau_i) = \begin{cases} 1 & \text{if } (\bar{n}_3, \bar{n}_2, \bar{n}_1) \in A, \\ -1 & \text{if } (\bar{n}_3, \bar{n}_2, \bar{n}_1) \in B, \end{cases}$$

where $A = \{(\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3), (\mathbf{e}_1, -\mathbf{e}_2, -\mathbf{e}_3)\}$ and $B = \{(\mathbf{e}_1, -\mathbf{e}_2, \mathbf{e}_3), (\mathbf{e}_1, \mathbf{e}_2, -\mathbf{e}_3)\}$. Therefore for each $i \in I_+^3$,

$$\epsilon(\tau_i) = \epsilon_{tm}(i) \epsilon_b(i). \quad (4.4)$$

Let $j \in I_-^3$, i.e., $D'_j \rightarrow D'_{j+1}$ is a move of type R_3 . Let τ'_j be the triple point in $M'_{(j)}$. Let n_1 , n_2 and n_3 be the co-orientations of the bottom, the middle and the top arcs of D'_j in $B'_{(j)}$, respectively. By an ambient isotopy,

we deform $M'_{(j)}$ to the standard form of the neighborhood of the triple point τ'_j . Let $\bar{n}_1$, $\bar{n}_2$, and $\bar{n}_3$ be the co-orientations corresponding to n_1 , n_2 , and n_3 , respectively. Without loss of generality, we may assume $\bar{n}_3 = \mathbf{e}_1$, $\bar{n}_2 = \epsilon \mathbf{e}_2$ and $\bar{n}_1 = \epsilon' \mathbf{e}_3$ for some $\epsilon, \epsilon' \in \{1, -1\}$.

Let c be the crossing between the top and the middle arcs in $B'_{(j)}$. It is easily seen that $\epsilon = \text{sign}(c)$ (cf. Figure 12). By (4.1), $\epsilon = \text{sign}(c) = \epsilon_{tm}(j)$. Hence $\bar{n}_2 = \epsilon_{tm}(j) \mathbf{e}_2$.

The sign ϵ' depends on the co-orientation n_1 of the bottom arc. If n_1 points from c , then $\epsilon' = -1$. If n_1 points toward c , then $\epsilon' = 1$. So, by (4.2), $\epsilon' = -\epsilon_b(j)$ and hence $\bar{n}_1 = -\epsilon_b(j) \mathbf{e}_3$.

On the other hand, by definition, $\epsilon(\tau'_j) = 1$ if $\bar{n}_3, \bar{n}_2$, and $\bar{n}_1$ in this order match the given (right-handed) orientation of $\mathbb{R}^3$. Otherwise, $\epsilon(\tau'_j) = -1$. This gives

$$\epsilon(\tau'_j) = \begin{cases} 1 & \text{if } (\bar{n}_3, \bar{n}_2, \bar{n}_1) \in B, \\ -1 & \text{if } (\bar{n}_3, \bar{n}_2, \bar{n}_1) \in A. \end{cases}$$

Therefore for each $j \in I_-^3$,

$$\epsilon(\tau'_j) = -\epsilon_{tm}(j) \epsilon_b(j). \quad (4.5)$$

We show that for each $i \in I_+^3$, $B_\theta(i, \mathcal{C}) = B_\theta(\tau_i, \mathcal{C})$ and that for each $j \in I_-^3$, $B_\theta(j, \mathcal{C}) = B_\theta(\tau'_j, \mathcal{C})^{-1}$.

Let $i \in I_+^3$ (or $j \in I_-^3$). There are two cases: The bottom arc meets the source region of the crossing c or not (see Figure 12). In this proof, we denote $M_{(i)}$ (or $M'_{(j)}$) by M and $[t_i, t_{i+1}]$ (or $[t'_{j+1}, t'_j]$) by I .

Case I: Consider $i \in I_+^3$ (or $j \in I_-^3$) such that the bottom arc in $B_{(i)}$ (or $B'_{(j)}$) hits the source region R of c .

The top (or the middle, resp.) sheet in M corresponds to the top (or the middle, resp.) arc times I . As shown in Figure 12, $R \times I$ is divided into two (3-dimensional) regions by the bottom sheet whose color is x_1 . One of them is the source region $\mathcal{R}$ of the triple point τ_i (or τ'_j). The colors of the top arc and the middle arc facing the source region R of c are the colors of the top and the middle sheets facing $\mathcal{R}$. From the equalities (4.4) and (4.5), we see that $B_\theta(i, \mathcal{C}) = \theta(x_1, x_2, x_3)^{\epsilon_{tm}(i) \epsilon_b(i)} = \theta(x_1, x_2, x_3)^{\epsilon(\tau_i)} = B_\theta(\tau_i, \mathcal{C})$ and $B_\theta(j, \mathcal{C}) = \theta(x_1, x_2, x_3)^{\epsilon_{tm}(j) \epsilon_b(j)} = \theta(x_1, x_2, x_3)^{-\epsilon(\tau'_j)} = B_\theta(\tau'_j, \mathcal{C})^{-1}$.

Case II: Consider $i \in I_+^3$ ($j \in I_-^3$) such that the bottom arc in $B_{(i)}$ (or $B'_{(j)}$) does not meet the source region R of c .

Similar to the case I, the second and third coordinates of $B_\theta(i, \mathcal{C})$ (or $B_\theta(j, \mathcal{C})$) are the same as those of $B_\theta(\tau_i, \mathcal{C})$ (or $B_\theta(\tau'_j, \mathcal{C})$). As illustrated in

Figure 12, $R' \times I$ is divided into two (3-dimensional) regions by the bottom sheet whose color is a , where a is the color of the bottom arc in R' . Since the co-orientation of the top sheet is from $R \times I$ to $R' \times I$, $a = x_1 * x_3$. Thus $x_1 = a * \overline{x_3}$. Therefore $B_\theta(i, \mathcal{C}) = \theta(x_1, x_2, x_3)^{\epsilon_{tm}(i)\epsilon_b(i)} = \theta(x_1, x_2, x_3)^{\epsilon(\tau_i)} = B_\theta(\tau_i, \mathcal{C})$ and $B_\theta(j, \mathcal{C}) = B_\theta(\tau'_j, \mathcal{C})^{-1}$.

This completes the proof of Claim and hence the proof of Theorem 4.5. $\square$

Example 4.6. We consider the oriented marked graph diagram 10_2 of the 2-twist spun trefoil $\mathcal{L}$ in Figure 15. Let

$$\theta = \chi_{0,1,0}^{-1} \chi_{0,2,0} \chi_{0,2,1}^{-1} \chi_{1,0,1} \chi_{1,0,2} \chi_{2,0,2} \chi_{2,1,2} \in Z_Q^3(R_3; \mathbb{Z}_3),$$

where $\chi_{x,y,z}(a,b,c) = u$ if $(x,y,z) = (a,b,c)$, $\chi_{x,y,z}(a,b,c) = 1$ otherwise, and $\mathbb{Z}_3 = \langle u \mid u^3 = 1 \rangle$ is the cyclic group of order 3. Consider sequences of link diagrams from the positive and negative resolutions to trivial link diagrams are as shown in Figures 13 and 14, respectively. Then $I_+^3 = \phi$ and $I_-^3 = \{2, 3, 4, 5, 8, 10\}$. The (Boltzman) weights are $B_\theta(2, \mathcal{C}) = \theta(y, y, x) = 1$, $B_\theta(3, \mathcal{C}) = \theta(x * y, x, x)^{-1} = 1$, $B_\theta(4, \mathcal{C}) = \theta(x, x * y, x)^{-1}$, $B_\theta(5, \mathcal{C}) = \theta(x, y, x)^{-1}$, $B_\theta(8, \mathcal{C}) = \theta(y, x, y)$ and $B_\theta(10, \mathcal{C}) = \theta(x * y, x, x * y)$ for $x, y \in R_3$. Therefore

$$\begin{aligned} \Phi_\theta(\mathcal{L}) &= \sum_{(x,y) \in R_3 \times R_3} \theta(x, x * y, x) \theta(x, y, x) \theta(y, x, y)^{-1} \theta(x * y, x, x * y)^{-1} \\ &= 3 + 6u. \end{aligned}$$

This matches the computation in [5].

For a surface-link $\mathcal{L}$, the *ch-index* $\chi(\mathcal{L})$ is defined by $\min_\Gamma \chi(\Gamma)$, where Γ is a marked graph diagram presenting $\mathcal{L}$ and $\chi(\Gamma)$ is the sum of the number of crossings of Γ and that of vertices of Γ .

Example 4.7. Let $\mathcal{L}$ be an oriented surface-link with $\chi(\mathcal{L}) \leq 10$ presented by a marked graph diagram in Figure 15 (see [32], for more details).

Let R_3 be the dihedral quandle of order 3 and θ the 3-cocycle in Example 4.6. Let S_4 be the tetrahedral quandle in Example 3.1 and let $\eta =$

$$\chi_{0,1,0} \chi_{0,2,1} \chi_{0,2,3} \chi_{0,3,0} \chi_{0,3,1} \chi_{0,3,2} \chi_{1,0,1} \chi_{1,0,3} \chi_{1,2,0} \chi_{1,3,1} \chi_{2,0,3} \chi_{2,1,0} \chi_{2,1,3} \chi_{2,3,2}$$

in $Z_Q^3(S_4; \mathbb{Z}_2)$, where $\chi_{x,y,z}(a,b,c) = t$ if $(x,y,z) = (a,b,c)$, $\chi_{x,y,z}(a,b,c) = 1$ otherwise, and $\mathbb{Z}_2 = \langle t \mid t^2 = 1 \rangle$. Then $\Phi_\theta(\mathcal{L})$ and $\Phi_\eta(\mathcal{L})$ are as in the table below.




Figure 13: A sequence of link diagrams for positive resolution of 10_2




Figure 14: A sequence of link diagrams for negative resolution of 10_2




Figure 15: Oriented marked graph diagrams Γ with $\chi(\Gamma) \leq 10$

$\mathcal{L}$	$\Phi_\theta(\mathcal{L})$	$\Phi_\eta(\mathcal{L})$	$\mathcal{L}$	$\Phi_\theta(\mathcal{L})$	$\Phi_\eta(\mathcal{L})$
0_1	3	4	10_2	$3+6u$	4
2_1^1	3	4	10_3	3	$4+12t$
$6_1^{0,1}$	3	4	10_1^1	9	16
8_1	9	16	$10_1^{0,1}$	3	4
$8_1^{1,1}$	3	4	$10_2^{0,1}$	3	4
9_1	9	16	$10_1^{1,1}$	3	4
$9_1^{0,1}$	3	4	$10_1^{0,0,1}$	9	16
10_1	3	4			

Table: $\Phi_\theta(\mathcal{L})$ and $\Phi_\eta(\mathcal{L})$ with $\chi(\mathcal{L}) \leq 10$

In [32], K. Yoshikawa introduced the notion of a marked graph diagram of *triangle type*. It is seen that the quandle cocycle invariant $\Phi_\theta(\mathcal{L})$ of an oriented surface-link $\mathcal{L}$ presented by a marked graph diagram of triangle type is equal to $\#\text{Col}_X(\mathcal{L})$ for any finite quandle X and any 3-cocycle $\theta \in Z_Q^3(X; A)$, where $\#\text{Col}_X(\mathcal{L})$ denotes the cardinality of the set $\text{Col}_X(\mathcal{L})$.

In [30], M. Soma gave an enumeration of surface-links presented by marked graph diagrams of *square type*; $A_n, B_n, C_n, D_n, E_n, F_n, G_n, H_n$ and I (See [22], [30, Theorems 1.1 and 1.2]). We remark that surface-links presented by marked graph diagrams A_n and B_n are orientable for all $n \geq 2$, and surface-links presented by marked graph diagrams E_n, F_n, G_n, H_n and I are also orientable for all odd integers $n \geq 3$. See Figure 16.

We observe that for any finite quandle X and $\theta \in Z_Q^3(X; A)$, the quandle cocycle invariant $\Phi_\theta(\mathcal{L})$ of an oriented surface-link $\mathcal{L}$ presented by a marked graph diagram of square type is equal to $\#\text{Col}_X(\mathcal{L})$ except for the surface-link presented by G_n .




Figure 16: Marked graph diagrams of square type

For an oriented surface-link $\mathcal{L}$, we denote the same surface-link as $\mathcal{L}$ but with the opposite orientations on all the components of $\mathcal{L}$ by $-\mathcal{L}$. An oriented surface-link $\mathcal{L}$ is said to be *invertible* if it is equivalent to $-\mathcal{L}$; otherwise *non-invertible*. The quandle cocycle invariant provides a diagrammatic method of detecting non-invertibility of surface-links (cf. [3, Section 3]).

Theorem 4.8. *For every integer $k \geq 0$, the oriented surface-links presented by marked graph diagrams G_{18k+3} and G_{18k+15} in Figure 16 are non-invertible.*

Proof. Let X be the dihedral quandle of order 3 and $\theta \in Z_Q^3(X; A)$ the 3-cocycle in Example 4.6. Let $\mathcal{L}$ be the oriented surface-link presented by the oriented marked graph diagram G_n in Figure 16. Then $\Phi_\theta(\mathcal{L}) = 3 + 6u$ and $\Phi_\theta(-\mathcal{L}) = 3 + 6u^2$ if $n = 18k + 3$, and $\Phi_\theta(\mathcal{L}) = 3 + 6u^2$ and $\Phi_\theta(-\mathcal{L}) = 3 + 6u$ if $n = 18k + 15$ for any integer $k \geq 0$. This shows that $\mathcal{L}$ and $-\mathcal{L}$ are not equivalent for any $k \geq 0$ and completes the proof. $\square$

On the other hand, it is shown that for every integer $m \geq 1$, the oriented surface-links presented by marked graph diagrams F_{2m+1} and H_{2m+1} in Figure 16 are all non-invertible [22, Theorem 7.4].

5 Shadow quandle cocycle invariants of oriented surface-links

In this section, we recall shadow quandle cocycle invariants of oriented surface-links (cf. [4]).

Let X be a quandle and let $\mathcal{B}$ be a broken surface diagram of an oriented surface-link $\mathcal{L}$. Let $S(\mathcal{B})$ be the set of sheets of $\mathcal{B}$ and $R(\mathcal{B})$ be the set of the complementary regions of $\mathcal{B}$ in $\mathbb{R}^3$. Let $\mathcal{C} : S(\mathcal{B}) \rightarrow X$ be a coloring of $\mathcal{B}$. A *shadow coloring* of $\mathcal{B}$ (extending a given coloring $\mathcal{C}$) is a map $\tilde{\mathcal{C}} : S(\mathcal{B}) \cup R(\mathcal{B}) \rightarrow X$ satisfying the conditions:

- The restriction of $\tilde{\mathcal{C}}$ to $S(\mathcal{B})$ is a given coloring $\mathcal{C}$.
- If two adjacent regions f_1 and f_2 are separated by a sheet e and the co-orientation of e points from f_1 to f_2 , then $\tilde{\mathcal{C}}(f_1) * \tilde{\mathcal{C}}(e) = \tilde{\mathcal{C}}(f_2)$.

Let $\text{Col}_X^S(\mathcal{B})$ be the set of all shadow colorings of $\mathcal{B}$ by X .

Let $\tilde{\mathcal{C}}$ be a shadow coloring of $\mathcal{B}$. Let τ be a triple point and let $\mathcal{R}$ be the source region of τ . Let $\theta \in Z_Q^4(X; A)$. Define the *shadow (Boltzman) weight* at τ by

$$B_\theta^S(\tau, \tilde{\mathcal{C}}) = \theta(y, x_1, x_2, x_3)^{\epsilon(\tau)},$$

where $\epsilon(\tau)$ is the sign of τ , y is the color of $\mathcal{R}$ and x_1, x_2 and x_3 are the colors of the bottom, the middle and the top sheets facing $\mathcal{R}$, respectively. See Figure 17 for $\epsilon(\tau) = 1$. The *shadow partition function* of $\mathcal{B}$ (associated to θ) is defined by

$$\Phi_\theta^s(\mathcal{B}) = \sum_{\tilde{\mathcal{C}} \in \text{Col}_X^S(\mathcal{B})} \prod_{\tau \in T(\mathcal{B})} B_\theta^S(\tau, \tilde{\mathcal{C}}) \in \mathbb{Z}[A].$$




Figure 17: Shadow (Boltzman) weight at τ with $\epsilon(\tau) = 1$

Theorem 5.1 ([4]). *Let $\mathcal{B}$ be a broken surface diagram of an oriented surface-link $\mathcal{L}$. The shadow partition function $\Phi_\theta^s(\mathcal{B})$ does not depend on the choice of a broken surface diagram. Thus it is an invariant of $\mathcal{L}$.*

We denote $\Phi_\theta^s(\mathcal{B})$ by $\Phi_\theta^s(\mathcal{L})$ and call it a *shadow quandle cocycle invariant* of $\mathcal{L}$ associated to $\theta \in Z_Q^4(X; A)$.

There is a generalized version of the shadow quandle cocycle invariant.

Let X be a quandle. The *associated group*, G_X , of X is $\langle x \in X; x * y = y^{-1}xy \ (x, y \in X) \rangle$. An *X-set* is a set Y equipped with a right action of the associated group G_X . We denote by $y * g$ the image of an element $y \in Y$ by the action $g \in G_X$.

Let X be a quandle and Y an X -set. For each positive integer n , let $C_n^R(X)_Y$ be the free abelian group generated by the elements $(y, x_1, \dots, x_n)$ where $y \in Y$ and $x_1, \dots, x_n \in X$. Let $C_0(X)_Y = \mathbb{Z}(Y)$, the free abelian group on Y , and let $C_n^R(X)_Y = \{0\}$ for $n < 0$. Define a homomorphism

$\partial_n : C_n^R(X)_Y \rightarrow C_{n-1}^R(X)_Y$ by

$$\begin{aligned} \partial_n(y, x_1, x_2, \dots, x_n) = & \sum_{i=1}^n (-1)^i [(y, x_1, x_2, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \\ & - (y * x_i, x_1 * x_i, x_2 * x_i, \dots, x_{i-1} * x_i, x_{i+1}, \dots, x_n)] \end{aligned}$$

for $n \geq 2$ and $\partial_n = 0$ for $n \leq 1$. Then $C_*^R(X)_Y = \{C_n^R(X)_Y, \partial_n\}$ is a chain complex. This chain complex is due to R. Fenn, C. Rourke and B. Sanderson ([8, 9]). Let $D_n^Q(X)_Y$ be the subgroup of $C_n^R(X)_Y$ generated by $(y, x_1, \dots, x_n)$ with $x_i = x_{i+1}$ for some $i \in \{1, \dots, n-1\}$ if $n \geq 2$; otherwise let $D_n^Q(X)_Y = \{0\}$. Then $C_*^D(X)_Y = \{D_n^Q(X)_Y, \partial_n\}$ is a subcomplex of $C_*^R(X)_Y$. Put $C_n^Q(X)_Y = C_n^R(X)_Y / D_n^Q(X)_Y$, and consider the quotient chain complex $C_*^Q(X)_Y = \{C_n^Q(X)_Y, \partial_n\}$. For an abelian group A , we define chain and cochain complexes by $C_*^Q(X, A)_Y = C_*^Q(X)_Y \otimes A$ and $C_Q^*(X, A)_Y = \text{Hom}(C_*^Q(X)_Y, A)$. The homology and cohomology groups are denoted by $H_*^Q(X, A)_Y$ and $H_Q^*(X, A)_Y$, respectively. For more details, see [17, 18].

Let X be a quandle, Y an X -set and let $\mathcal{B}$ be a broken surface diagram of an oriented surface-link $\mathcal{L}$. Let $S(\mathcal{B})$ be the set of sheets of $\mathcal{B}$ and $R(\mathcal{B})$ the set of the complementary regions of $\mathcal{B}$ in $\mathbb{R}^3$. Let $\mathcal{C} : S(\mathcal{B}) \rightarrow X$ be a coloring of $\mathcal{B}$. A *shadow coloring* of $\mathcal{B}$ (extending a given coloring $\mathcal{C}$) by (X, Y) is a map $\tilde{\mathcal{C}} : S(\mathcal{B}) \cup R(\mathcal{B}) \rightarrow X \cup Y$ satisfying the conditions:

- $\tilde{\mathcal{C}}(S(\mathcal{B})) \subset X$ and $\tilde{\mathcal{C}}(R(\mathcal{B})) \subset Y$.
- The restriction of $\tilde{\mathcal{C}}$ to $S(\mathcal{B})$ is a given coloring $\mathcal{C}$.
- If two adjacent regions f_1 and f_2 are separated by a sheet e and the co-orientation of e points from f_1 to f_2 , then $\tilde{\mathcal{C}}(f_1) * \tilde{\mathcal{C}}(e) = \tilde{\mathcal{C}}(f_2)$.

We denote by $\text{Col}_{(X,Y)}^S(\mathcal{B})$ the set of all shadow colorings of $\mathcal{B}$ by (X, Y) .

Proposition 5.2 (cf. [4]). *If $\mathcal{B}$ and $\mathcal{B}'$ present equivalent oriented surface-links, then there is a bijection between $\text{Col}_X^S(\mathcal{B})$ and $\text{Col}_X^S(\mathcal{B}')$, and there is a bijection between $\text{Col}_{(X,Y)}^S(\mathcal{B})$ and $\text{Col}_{(X,Y)}^S(\mathcal{B}')$.*

Let $\tilde{\mathcal{C}}$ be a shadow coloring of a broken surface diagram $\mathcal{B}$ by (X, Y) . Let τ be a triple point and let $\mathcal{R}$ be the source region of τ . Let $\theta \in Z_Q^3(X; A)_Y$. Define the *shadow (Boltzman) weight* at τ by

$$B_\theta^S(\tau, \tilde{\mathcal{C}}) = \theta(y, x_1, x_2, x_3)^{\epsilon(\tau)},$$

where $\epsilon(\tau)$ is the sign of τ , y is the color of $\mathcal{R}$ and x_1, x_2 and x_3 are the colors of the bottom, the middle and the top sheets facing $\mathcal{R}$, respectively. See Figure 17. The *shadow partition function* of $\mathcal{B}$ (associated to θ) is defined by

$$\Phi_\theta^s(\mathcal{B}) = \sum_{\tilde{\mathcal{C}} \in \text{Col}_{(X,Y)}^S(\mathcal{B})} \prod_{\tau \in T(\mathcal{B})} B_\theta^S(\tau, \tilde{\mathcal{C}}) \in \mathbb{Z}[A].$$

Theorem 5.3 (cf. [4]). *Let $\mathcal{B}$ be a broken surface diagram of an oriented surface-link $\mathcal{L}$. The shadow partition function $\Phi_\theta^s(\mathcal{B})$ does not depend on the choice of a broken surface diagram. Thus it is an invariant of $\mathcal{L}$.*

We denote $\Phi_\theta^s(\mathcal{B})$ by $\Phi_\theta^s(\mathcal{L})$ and call it a *shadow quandle cocycle invariant* of $\mathcal{L}$ associated to $\theta \in Z_Q^3(X; A)_Y$.

6 How to compute shadow quandle cocycle invariants from marked graph diagrams

In this section we give a method of computing shadow quandle cocycle invariants from marked graph diagrams.

Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$. Let $A(\Gamma)$ be the set of arcs of Γ and $R(\Gamma)$ the set of complementary regions of Γ in $\mathbb{R}^2$. Let X be a quandle and let Y be an X -set. Let $\mathcal{C} : A(\Gamma) \rightarrow X$ be a coloring of Γ by a quandle X . A *shadow coloring* of Γ (extending a given coloring $\mathcal{C}$) by X (or by (X, Y) , resp.) is a map $\tilde{\mathcal{C}} : A(\Gamma) \cup R(\Gamma) \rightarrow X$ (or a map $\tilde{\mathcal{C}} : A(\Gamma) \cup R(\Gamma) \rightarrow X \cup Y$, resp.) satisfying the conditions (2) and (3) (or the conditions (1)–(3), resp.):

- (1) $\tilde{\mathcal{C}}(A(\Gamma)) \subset X$ and $\tilde{\mathcal{C}}(R(\Gamma)) \subset Y$.
- (2) The restriction of $\tilde{\mathcal{C}}$ to $A(\Gamma)$ is a given coloring $\mathcal{C}$.
- (3) If two adjacent regions f_1 and f_2 are separated by an arc $e \in A(\Gamma)$ and the co-orientation of e points from f_1 to f_2 , then $\tilde{\mathcal{C}}(f_1) * \tilde{\mathcal{C}}(e) = \tilde{\mathcal{C}}(f_2)$.

Let $\text{Col}_X^S(\Gamma)$ (or $\text{Col}_{(X,Y)}^S(\Gamma)$, resp.) denote the set of all shadow colorings of Γ by X (or by (X, Y) , resp.).

Theorem 6.1. *Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$ and $\mathcal{B} = \mathcal{B}(\Gamma)$ an associated broken surface diagram of Γ . There is a bijection from $\text{Col}_X^S(\Gamma)$ to $\text{Col}_X^S(\mathcal{B})$, and a bijection from $\text{Col}_{(X,Y)}^S(\Gamma)$ to $\text{Col}_{(X,Y)}^S(\mathcal{B})$.*

Proof. Consider a shadow coloring of $\mathcal{B}$. The 0-level cross-section with the colors induced by the shadow coloring of $\mathcal{B}$ is a shadow coloring of Γ . By the same argument as in [1], we see that this gives a bijection from $\text{Col}_X^S(\Gamma)$ to $\text{Col}_X^S(\mathcal{B})$ and a bijection $\text{Col}_{(X,Y)}^S(\Gamma)$ to $\text{Col}_{(X,Y)}^S(\mathcal{B})$. $\square$

Corollary 6.2. *If Γ and Γ' present equivalent oriented surface-links, then there is a bijection from $\text{Col}_X^S(\Gamma)$ to $\text{Col}_X^S(\Gamma')$, and there is a bijection from $\text{Col}_{(X,Y)}^S(\Gamma)$ to $\text{Col}_{(X,Y)}^S(\Gamma')$.*

Proof. Let $\mathcal{B}(\Gamma)$ and $\mathcal{B}(\Gamma')$ be broken surface diagrams associated to Γ and Γ' , respectively. By Proposition 5.2 and Theorem 6.1, we see the result. $\square$

Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$ and let $\Gamma_+ = D_1 \rightarrow D_2 \rightarrow \cdots \rightarrow D_r = O$, $\Gamma_- = D'_1 \rightarrow D'_2 \rightarrow \cdots \rightarrow D'_s = O'$, ϵ_{tm} and ϵ_b be as in Section 4. Let $\tilde{\mathcal{C}} : A(\Gamma) \cup R(\Gamma) \rightarrow X$ or $\tilde{\mathcal{C}} : A(\Gamma) \cup R(\Gamma) \rightarrow X \cup Y$ be a shadow coloring of Γ . Let $i \in I_+^3$ (or $j \in I_-^3$). Let R be the source region of the crossing c between the top arc and the middle arc in $D_i \cap B_{(i)}$ (or $D'_j \cap B'_{(j)}$). Let R' be the opposite region of R with respect to the top arc.

Let x_1 , x_2 and x_3 be as in Section 4. There are two cases, the bottom arc intersects with the source region R or not. If not, we consider two cases, $\epsilon_b(i) = 1$ or $\epsilon_b(i) = -1$ (See Figure 18). In the case where the bottom arc hits the source region R , the region R is divided by the bottom arc. Let y be the color of the divided region of R such that the co-orientation of the bottom arc points from that region. In the case where the bottom arc does not intersect with the source region R and $\epsilon_b(i) = 1$, let y be the element $s = \tilde{\mathcal{C}}(R)$. In the case where the bottom arc does not meet the source region R and $\epsilon_b(i) = -1$, let y be the element $s * \overline{x_1}$, where $s = \tilde{\mathcal{C}}(R)$. For $j \in I_-^3$, let y be the element defined in the same way with $i \in I_+^3$.

Definition 6.3. Let $\theta \in Z_Q^4(X; A)$ be a 4-cocycle or let $\theta \in Z_Q^3(X; A)_Y$ be a 3-cocycle. The *shadow (Boltzman) weight* for $i \in I_+^3 \amalg I_-^3$ is defined by

$$B_\theta^S(i, \tilde{\mathcal{C}}) = \theta(y, x_1, x_2, x_3)^{\epsilon_{tm}(i)\epsilon_b(i)}.$$




Figure 18: The triple point τ_i

Definition 6.4. Let Γ be a marked graph diagram of an oriented surface-link $\mathcal{L}$. The *shadow partition function* of Γ (associated to θ) is defined by

$$\Phi_{\theta}^s(\Gamma) = \sum_{\tilde{C}} \left(\prod_{i \in I_{+}^3} B_{\theta}^S(i, \tilde{C}) \prod_{j \in I_{-}^3} B_{\theta}^S(j, \tilde{C})^{-1} \right),$$

where $\tilde{C}$ runs all shadow colorings of Γ by X when $\theta \in Z_Q^4(X; A)$ or all shadow colorings of Γ by (X, Y) when $\theta \in Z_Q^3(X; A)_Y$.

Theorem 6.5. *Let $\mathcal{L}$ be an oriented surface-link and let Γ be a marked graph diagram presenting $\mathcal{L}$. For any $\theta \in Z_Q^4(X; A)$ or $\theta \in Z_Q^3(X; A)_Y$, $\Phi_\theta^s(\mathcal{L}) = \Phi_\theta^s(\Gamma)$.*

Proof. Let $\mathcal{B} = \mathcal{B}(\Gamma)$ be a broken surface diagram associated to Γ . It suffices to show that $\Phi_\theta^s(\mathcal{B}) = \Phi_\theta^s(\Gamma)$.

We define $M_{(i)}$ and $M'_{(j)}$ for any $i \in I_+^3$ and $j \in I_-^3$ as in the proof of Theorem 4.5. We have that $T(\mathcal{B}) = \{\tau_i \mid i \in I_+^3\} \cup \{\tau'_j \mid j \in I_-^3\}$, where τ_i is the triple point in $M_{(i)}$ for $i \in I_+^3$ and τ'_j is the triple point in $M'_{(j)}$ for $j \in I_-^3$.

Let $\tilde{\mathcal{C}}$ be a shadow coloring of $\mathcal{B}$ by X when $\theta \in Z_Q^4(X; A)$ or a shadow coloring of $\mathcal{B}$ by (X, Y) when $\theta \in Z_Q^3(X; A)_Y$, and let $\mathcal{C} \in \text{Col}_X(\mathcal{B})$ be the restriction of $\tilde{\mathcal{C}}$ to the set $S(\mathcal{B})$.

We show that $B_\theta^S(i, \tilde{\mathcal{C}}) = B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ for each $i \in I_+^3$.

The exponents appearing in $B_\theta^S(i, \tilde{\mathcal{C}})$ and $B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ are identical. The second, the third and the fourth coordinates of $B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ are the same as $B_\theta(\tau, \mathcal{C})$ for every triple point τ . Also, the second, third and fourth coordinates of $B_\theta^S(i, \tilde{\mathcal{C}})$ are the same as $B_\theta(i, \mathcal{C})$ for each $i \in I_+^3$. Combining these facts, the second, the third and the fourth coordinates of $B_\theta^S(i, \tilde{\mathcal{C}})$ are the same as those of $B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ for any $i \in I_+^3$.

It remains to show that the first coordinate of $B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ is identical with that of $B_\theta^S(i, \tilde{\mathcal{C}})$ for any $i \in I_+^3$. The first coordinate of $B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ is the color of the source region $\mathcal{R}$ of the triple point τ_i .

Case I : The bottom arc intersects with the source region R of the crossing between the top and middle arc in $B_{(i)}$.

The top (or the middle, resp.) sheet corresponds to the top (or the middle, resp.) arc times $[t_i, t_{i+1}]$. Also, the quadrant between the top and middle sheets with the co-orientations outward is divided into two (3-dimensional) regions by a bottom sheet whose color is same as that of the bottom arc in R . Therefore, the first coordinate of $B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ is $\tilde{\mathcal{C}}(R)$, where R is the source region of the crossing c between the top and middle arcs in $B_{(i)}$. Therefore $B_\theta^S(i, \tilde{\mathcal{C}}) = B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ for all $i \in I_+^3$.

Case II : The bottom arc does not hit the source region R of the crossing between the top and middle arc in $B_{(i)}$.

Let $\tilde{\mathcal{C}}(R) = s$. The quadrant corresponding to $R \times [t_i, t_{i+1}]$ is divided into two (3-dimensional) regions by a bottom sheet whose color is $x_1 = a * \overline{x_3}$. If $\epsilon_b(i) = 1$, then the co-orientation of the bottom sheet in that quadrant is from the region which has a color s (see the case II-1 in Figure 18). Therefore the color y of the source region $\mathcal{R}$ of the triple point τ_i is s . Otherwise, the

co-orientation of the bottom sheet in that quadrant points to the region whose color is s (see the case II-2 in Figure 18). In addition, the color of the bottom sheet in that quadrant is x_1 . Thus the color y of $\mathcal{R}$ is $s * \overline{x_1}$. Therefore $B_\theta^S(i, \tilde{\mathcal{C}}) = B_\theta^S(\tau_i, \tilde{\mathcal{C}})$ for all $i \in I_+^3$.

For $j \in I_-^3$, it is similarly seen that $B_\theta^S(j, \tilde{\mathcal{C}}) = B_\theta^S(\tau'_j, \tilde{\mathcal{C}})^{-1}$.

Hence we have $\Phi_\theta^s(\mathcal{B}) = \Phi_\theta^s(\Gamma)$ for all $j \in I_-^3$. $\square$

7 Symmetric quandle cocycle invariants of unoriented surface-links

This section is devoted to recalling symmetric quandle cocycle invariants of unoriented surface-links (cf. [17, 18]).

Let X be a quandle. A map $\rho : X \rightarrow X$ is a *good involution* if it is an involution (i.e., $\rho \circ \rho = \text{id}$) such that $\rho(x * y) = \rho(x) * y$ and $x * \rho(y) = x * \bar{y}$ for any $x, y \in X$. Such a pair (X, ρ) is called a *quandle with a good involution* or a *symmetric quandle*.

Example 7.1. ([17, 18]) Let G be a group. The inversion, $\text{inv}(G) : G \rightarrow G; g \mapsto g^{-1}$, is a good involution of $\text{conj}(G)$. We call $(\text{conj}(G), \text{inv}(G))$ the *conjugation symmetric quandle*.

The *associated group*, $G_{(X, \rho)}$, of a symmetric quandle (X, ρ) is $G_{(X, \rho)} = \langle x \in X; x * y = y^{-1}xy \ (x, y \in X), \ \rho(x) = x^{-1} \ (x \in X) \rangle$. An (X, ρ) -*set* is a set Y equipped with a right action of the associated group $G_{(X, \rho)}$. We denote by $y * g$ the image of an element $y \in Y$ by the action $g \in G_{(X, \rho)}$.

Let (X, ρ) be a symmetric quandle and Y an (X, ρ) -set. Let $C_*^R(X)_Y = \{C_n^R(X)_Y, \partial_n\}$ be the chain complex of X with Y , and $C_*^D(X)_Y = \{D_n^Q(X)_Y, \partial_n\}$ be the sub-complex of $C_*^R(X)_Y$ as in Section 5.

Let $D_n^\rho(X)_Y$ be the subgroup of $C_n^R(X)_Y$ generated by

$$(y, x_1, \dots, x_n) + (y * x_j, x_1 * x_j, \dots, x_{j-1} * x_j, \rho(x_j), x_{j+1}, \dots, x_n)$$

for $j \in 1, \dots, n$ if $n \geq 2$; otherwise let $D_n^\rho(X)_Y = \{0\}$.

Define $C_n^{Q, \rho}(X)_Y$ to be $C_n^R(X)_Y / (D_n^Q(X)_Y + D_n^\rho(X)_Y)$, and we have the quotient complex $C_*^{Q, \rho}(X)_Y = \{C_n^{Q, \rho}(X)_Y, \partial_n\}$. For an abelian group A , we define chain and cochain complexes by $C_*^{Q, \rho}(X, A)_Y = C_*^{Q, \rho}(X)_Y \otimes A$ and $C_{Q, \rho}^*(X, A)_Y = \text{Hom}(C_*^{Q, \rho}(X)_Y, A)$, respectively. The homology and cohomology groups are denoted by $H_*^{Q, \rho}(X, A)_Y$ and $H_{Q, \rho}^*(X, A)_Y$, respectively. For details, see [17, 18].

Let $\mathcal{B}$ be an unoriented broken surface diagram. When we divide over-sheets at the double curves, we call the sheets of the result *semi-sheets* of $\mathcal{B}$. Each semi-sheet is a compact orientable surface in $\mathbb{R}^3$ (cf. [16]).

Consider an assignment of normal orientation and an element of X to each semi-sheet of $\mathcal{B}$. A *basic inversion* is an operation which reverses the normal orientation of a semi-sheet and changes the element x assigned to the semi-sheet by $\rho(x)$. See Figure 19.



Figure 19: A basic inversion

We would rather use the terminology ‘normal orientation’ than ‘co-orientation’ when $\mathcal{B}$ is an unoriented broken surface diagram.

An (X, ρ) -coloring of $\mathcal{B}$ is the equivalence class of an assignment of a normal orientation and an element of X to each semi-sheet of $\mathcal{B}$ satisfying the coloring condition below. Here the equivalence relation is generated by basic inversions.

- By basic inversions, assume the normal orientations of semi-sheets around a double point curve to be as in Figure 20. Then $x_1 * x_3 = x_2$ and $x_3 = x_4$.




Figure 20: Coloring conditions

Let Y be an (X, ρ) -set. An $(X, \rho)_Y$ -coloring of $\mathcal{B}$ is an (X, ρ) -coloring of $\mathcal{B}$ with an assignment of an element of Y to each complementary region of $\mathcal{B}$ satisfying the following condition.

- Suppose that adjacent regions f_1 and f_2 separated by a semi-sheet e are labeled by y_1 and y_2 . If the semi-sheet e is labeled by x and the normal orientation of e points from f_1 to f_2 , then $y_1 * x = y_2$.

Proposition 7.2 ([17, 18]). *Let (X, ρ) be a symmetric quandle and Y an (X, ρ) -set. If two broken surface diagrams present equivalent unoriented surface-links, then there is a bijection between the sets of (X, ρ) -colorings of the broken surface diagrams, and there is a bijection between the sets of $(X, \rho)_Y$ -colorings of them.*

Let $\mathcal{B}$ be an unoriented broken surface diagram. Fix an $(X, \rho)_Y$ -coloring of $\mathcal{B}$, say $\tilde{\mathcal{C}}$. For a triple point τ of $\mathcal{B}$, there are eight complementary regions of $\mathcal{B}$ around τ (Some of them may be the same). Choose one of them, say f , which we call a *specified region* for τ , and let y be the label of f .

Let e_1 , e_2 and e_3 be the bottom semi-sheet, the middle semi-sheet and the top semi-sheet at τ , respectively, which face the region f . By basic inversions, we assume that the normal orientations n_1 , n_2 and n_3 of them point from f to the opposite regions. Let x_1 , x_2 and x_3 be the labels of them, respectively. The *sign* of τ with respect to the region f is $+1$ (or -1) if the triple of normal orientations (n_3, n_2, n_1) does (or does not) match the orientation of $\mathbb{R}^3$. Let $\theta \in Z_{Q, \rho}^3(X, A)_Y$. The *symmetric (Boltzman) weight* $B_\theta^{Sym}(\tau, \tilde{\mathcal{C}})$ of τ is defined to be

$$B_\theta^{Sym}(\tau, \tilde{\mathcal{C}}) = \theta(y, x_1, x_2, x_3)^{\epsilon(\tau)},$$

where $\epsilon(\tau)$ is the sign of τ . See Figure 21.




Figure 21: Symmetric Boltzman weights

The *symmetric partition function* of $\mathcal{B}$ (associated to θ) is defined by

$$\Phi_\theta^{Sym}(\mathcal{B}) = \sum_{\tilde{\mathcal{C}}} \prod_{\tau \in T(\mathcal{B})} B_\theta^{Sym}(\tau, \tilde{\mathcal{C}}) \in \mathbb{Z}[A],$$

where the sum is taken over all possible $(X, \rho)_Y$ -colorings $\tilde{\mathcal{C}}$ of $\mathcal{B}$. (The value of $B_\theta^{Sym}(\tau, \tilde{\mathcal{C}})$ is in the coefficient group A written multiplicatively).

Theorem 7.3 ([17, 18]). *Let $\mathcal{B}$ be a broken surface diagram of an unoriented surface-link $\mathcal{L}$. The symmetric partition function $\Phi_\theta^{Sym}(\mathcal{B})$ is an invariant of the unoriented surface-link $\mathcal{L}$.*

We denote $\Phi_\theta^{Sym}(\mathcal{B})$ by $\Phi_\theta^{Sym}(\mathcal{L})$ and call it the *symmetric quandle cocycle invariant* of $\mathcal{L}$ associated to θ .

8 How to compute symmetric quandle cocycle invariants from marked graph diagrams

Let Γ be a marked graph diagram of an unoriented surface-link $\mathcal{L}$ and let (X, ρ) be a symmetric quandle.

A *semi-arc* of Γ is a connected component of $\Gamma \setminus (C(\Gamma) \cup V(\Gamma))$, where $C(\Gamma)$ is the set of crossings and $V(\Gamma)$ is the set of marked vertices of Γ .

A *basic inversion* is an operation which reverses the normal orientation of a semi-arc and changes the element x assigned to the semi-arc by $\rho(x)$. See Figure 22.




Figure 22: A basic inversion

We say that an assignment of a normal orientation and an element of X to each semi-arc of Γ satisfies the *coloring conditions* if it satisfies the following conditions.

- For each marked vertex, using basic inversions, we assume that normal orientations of semi-arcs are as in Figure 23. Then $x_1 = x_2$.
- For each crossing, using basic inversions, we assume that normal orientations of semi-arcs are as in Figure 23. Then $x_1 * x_3 = x_2$ and $x_3 = x_4$.

An (X, ρ) -coloring of Γ is the equivalence class of an assignment of a normal orientation and an element of X to each semi-arc of Γ satisfying the coloring conditions. Here the equivalence relation is generated by basic inversions.




Figure 23: Coloring conditions

Let Y be an (X, ρ) -set. An $(X, \rho)_Y$ -coloring of Γ is an (X, ρ) -coloring with an assignment of an element of Y to each complementary region of Γ satisfying the following condition.

- Suppose that two adjacent regions f_1 and f_2 separated by a semi-arc e are labeled by y_1 and y_2 . If the semi-arc e is labeled by x and the normal orientation of e points from f_1 to f_2 , then $y_1 * x = y_2$.

Theorem 8.1. *Let (X, ρ) be a symmetric quandle and let Y be an (X, ρ) -set. Let Γ be an admissible marked graph diagram, and let $\mathcal{B} = \mathcal{B}(\Gamma)$ be a broken surface diagram associated with Γ . There is a bijection from the set of $(X, \rho)_Y$ -colorings of $\mathcal{B}$ to that of Γ .*

Proof. By the same argument as in the proof of Theorem 6.1, we see the result. $\square$

Let Γ be an admissible marked graph diagram. Fix an $(X, \rho)_Y$ -coloring of Γ , say $\tilde{C}$. Then both resolutions Γ_+ and Γ_- have induced colorings.




Figure 24: Induced colorings on Γ_+ and Γ_-

Let $\Gamma_+ = D_1 \rightarrow D_2 \rightarrow \cdots \rightarrow D_r = O$ and $\Gamma_- = D'_1 \rightarrow D'_2 \rightarrow \cdots \rightarrow D'_s = O'$ be sequences of link diagrams as before. Let $i \in I_+^3 \amalg I_-^3$ and f the

complementary region of D_i in $B_{(i)}$ or $B'_{(i)}$ such that f does not intersect with the boundary $\partial B_{(i)}$ or $\partial B'_{(i)}$, respectively. Let e_1 , e_2 and e_3 be the bottom, the middle and the top semi-arcs facing the region f , respectively. By basic inversions, we assume that the normal orientations n_1 , n_2 and n_3 of them point outwards. Let x_1 , x_2 and x_3 be the labels of them, respectively. Define $\epsilon_{tm}(i) = 1$ if (n_3, n_2) matches with the given (right-handed) orientation of $\mathbb{R}^2$ and -1 otherwise. For a given 3-cocycle $\theta \in Z_{Q,\rho}^3(X, A)_Y$, we define the *symmetric (Boltzman) weight* at i to be

$$B_\theta^{Sym}(i, \tilde{\mathcal{C}}) = \theta(y, x_1, x_2, x_3)^{\epsilon_{tm}(i)}.$$




Figure 25: The symmetric (Boltzman) weight at $i \in I_+^3 \amalg I_-^3$

For a marked graph diagram Γ and an $(X, \rho)_Y$ -coloring $\tilde{\mathcal{C}}$, we define the *symmetric partition function* by

$$\Phi_\theta^{Sym}(\Gamma) = \sum_{\tilde{\mathcal{C}}} \left(\prod_{i \in I_+^3} B_\theta^{Sym}(i, \tilde{\mathcal{C}}) \prod_{j \in I_-^3} B_\theta^{Sym}(j, \tilde{\mathcal{C}})^{-1} \right),$$

where $\tilde{\mathcal{C}}$ runs over all $(X, \rho)_Y$ -colorings of Γ .

Theorem 8.2. *Let $\mathcal{L}$ be an unoriented surface-link and let Γ be a marked graph diagram presenting $\mathcal{L}$. For any 3-cocycle $\theta \in Z_{Q,\rho}^3(X, A)_Y$, the symmetric partition functions $\Phi_\theta^{Sym}(\Gamma)$ is equal to $\Phi_\theta^{Sym}(\mathcal{L})$.*

Proof. The proof of this theorem is similar to that of Theorem 6.5. Let $\mathcal{B} = \mathcal{B}(\Gamma)$ be the broken surface diagram associated to Γ , and let $\tilde{\mathcal{C}} \in \text{Col}_\theta^{Sym}(\mathcal{B})$ be an $(X, \rho)_Y$ -coloring of $\mathcal{B}$. We denote by the same symbol $\tilde{\mathcal{C}}$ for the corresponding $(X, \rho)_Y$ -coloring of Γ . As the oriented case, the set of triple points is $T(\mathcal{B}) = \{\tau_i \mid i \in I_+^3\} \cup \{\tau'_j \mid j \in I_-^3\}$, where τ_i is the triple point in $M_{(i)}$ for $i \in I_+^3$ and τ'_j is the triple point in $M'_{(j)}$ for $j \in I_-^3$. Let $i \in I_+^3$. Since

we choose the normal orientation of the bottom arc such that $\epsilon_b(i) = 1$, we have $\epsilon(\tau_i) = \epsilon_{tm}(i)$. Thus $B_\theta^{Sym}(i, \tilde{\mathcal{C}}) = B_\theta^{Sym}(\tau_i, \tilde{\mathcal{C}})$. Similarly, for $j \in I_-^3$, we have $B_\theta^{Sym}(j, \tilde{\mathcal{C}}) = B_\theta^{Sym}(\tau'_j, \tilde{\mathcal{C}})^{-1}$. Hence we have $\Phi_\theta^{Sym}(\Gamma) = \Phi_\theta^{Sym}(\mathcal{B})$.




Figure 26: A triple point

□

Example 8.3. Let Γ be the unorientable marked graph diagram in Figure 27 representing two component $\mathbb{RP}^2$ -link $\mathcal{L}$ (Γ is a marked graph diagram $8_1^{-1, -1}$ in Yoshikawa table [32]).

Let X be the dihedral quandle of order 4, in which we rename the elements 0, 1, 2, 3 by e_1, e_2, e'_1, e'_2 , respectively. Let $\rho : X \rightarrow X$ be the antipodal




Figure 27: A symmetric coloring for Γ

map, i.e., $\rho(e_i) = e'_i$ ($i = 1, 2$). Let $Y = \{e\}$, which is an (X, ρ) -set. Let

$$\begin{aligned} \theta = & \chi_{e, e_1, e_2, e_1} \chi_{e, e'_1, e'_2, e_1} \chi_{e, e'_1, e_2, e'_1} \chi_{e, e_1, e'_2, e'_1} \\ & \chi_{e, e'_1, e_2, e_1}^{-1} \chi_{e, e_1, e'_2, e_1}^{-1} \chi_{e, e_1, e_2, e'_1}^{-1} \chi_{e, e'_1, e'_2, e'_1}^{-1} \\ & \chi_{e, e_2, e_1, e_2}^{-1} \chi_{e, e'_2, e'_1, e_2}^{-1} \chi_{e, e'_2, e_1, e'_2}^{-1} \chi_{e, e_2, e'_1, e'_2}^{-1} \\ & \chi_{e, e'_2, e_1, e_2} \chi_{e, e_2, e'_1, e_2} \chi_{e, e_2, e_1, e'_2} \chi_{e, e'_2, e'_1, e'_2} \in Z_Q^3(X; \mathbb{Z}), \end{aligned}$$

where $\chi_{x, y, z, w}(a, b, c, d) = t$ if $(x, y, z, w) = (a, b, c, d)$, $\chi_{x, y, z, w}(a, b, c, d) = 1$ otherwise, and $\mathbb{Z} = \langle t \rangle$ is the infinite cyclic group (cf. [18, Example 9.3]).

Consider sequences of link diagrams from the positive and negative resolutions to trivial link diagrams as in Figures 28 and 29, respectively. From those figures, we get $I_+^3 = \{2, 3, 4, 6\}$ and $I_-^3 = \emptyset$. The symmetric (Boltzmann) weights are $B_\theta^{Sym}(2, \mathcal{C}) = \theta(e, y, \rho(y), \rho(x)) = \theta(e, y, y, \rho(x))^{-1} = 1$, $B_\theta^{Sym}(3, \mathcal{C}) = \theta(e, y, \rho(x), \rho(x)) = 1$, $B_\theta^{Sym}(4, \mathcal{C}) = \theta(e, x, y, \rho(x))$, $B_\theta^{Sym}(6, \mathcal{C}) = \theta(e, \rho(y), x, y)^{-1}$ for $(x, y) \in E$, where $E = \{(e_1, e_2), (e_1, e'_2), (e'_1, e_2), (e'_1, e'_2), (e_2, e_1), (e_2, e'_1), (e'_2, e_1), (e'_2, e'_1)\}$. Therefore

$$\begin{aligned} \Phi_\theta(\mathcal{L}) &= \sum_{\tilde{\mathcal{C}}} \left(\prod_{i \in I_+^3} B_\theta^{Sym}(i, \tilde{\mathcal{C}}) \prod_{j \in I_-^3} B_\theta^{Sym}(j, \tilde{\mathcal{C}})^{-1} \right) \\ &= \theta(e, x, y, \rho(x)) \theta(e, \rho(y), x, y)^{-1} \\ &= 4 + 2t^2 + 2t^{-2}. \end{aligned}$$

Acknowledgements. The first author was supported by JSPS KAKENHI Grant Number 26287013. The third author was supported by Basic Science Research Program through the National Research Foundation of Korea(NRF) funded by the Ministry of Education, Science and Technology (2013R1A1A2012446).




Figure 28: A sequence of link diagrams for Γ_+




Figure 29: A sequence of link diagrams for Γ_-

References

- [1] S. Ashihara, Calculating the fundamental biquandles of surface-links from their ch-diagrams. *J. Knot Theory Ramifications* **21** (2012), no. 10, 1250102 (23 pages).
- [2] J. S. Carter and M. Saito, *Knotted surfaces and their diagrams*, American Mathematical Society, 1998.
- [3] J. S. Carter, S. Kamada and M. Saito, Surfaces in 4-space, *Springer*, 2004.
- [4] J. S. Carter, S. Kamada and M. Saito, Geometric interpretations of quandle homology, *J. Knot Theory Ramifications* **10** (2001), no. 3, 345–386.
- [5] J. S. Carter, D. Jelsovsky, S. Kamada, L. Langford and M. Saito, Quandle cohomology and state-sum invariants of knotted curves and surfaces, *Trans. Amer. Math. Soc.* **355** (2003), 3947–3989.
- [6] J. Scott Carter, Daniel Jelsovsky, Seiichi Kamada and Masahico Saito, Computations of quandle cocycle invariants of knotted curves and surfaces, *Adv. in Math.* **157** (2001), no. 1, 36–94.
- [7] J. Scott Carter, Daniel Jelsovsky, Seiichi Kamada and Masahico Saito, Quandle homology groups, their Betti numbers, and virtual knots, *J. Pure Appl. Algebra*, **157** (2001), no. 2, 135–155.
- [8] R. Fenn, C. Rourke and B. Sanderson, Trunks and classifying spaces, *Appl. Categ. Structures* **3** (1995), no. 4, 321–356.
- [9] R. Fenn, C. Rourke and B. Sanderson, James bundles and applications, preprint (1996), <http://www.maths.warwick.ac.uk/cpr/ftp/james.ps>.
- [10] J. Flower, Cyclic Bordism and Rack Spaces, Ph.D. Dissertation, Warwick, 1995.
- [11] R. H. Fox, A quick trip through knot theory, in *Topology of 3-manifolds and Related Topics*, Prentice-Hall, Inc., Englewood Cliffs, N.J., 1962, 120–167.
- [12] M. T. Greene, Some Results in Geometric Topology and Geometry, Ph.D. Dissertation, Warwick, 1997.

- [13] Y. Joung, J. Kim and S. Y. Lee, Ideal coset invariants for surface-links in $\mathbb{R}^4$, *J. Knot Theory Ramifications* **22** (2013), no. 9, 1350052 (25 pages).
- [14] Y. Joung, S. Kamada and S. Y. Lee, Applying Lipson's state models to marked graph diagrams of surface-links, preprint (2014), arXiv:1411.5740 [math.GT]
- [15] D. Joyce, A classifying invariant of knots, the knot quandle, *J. Pure Appl. Algebra* **23** (1982), 37–65.
- [16] S. Kamada, Wirtinger presentations for higher dimensional manifold knots obtained from diagrams, *Fund. Math.* **168** (2001), 105–112.
- [17] S. Kamada, Quandles with good involutions, their homologies and knot invariants, in: *Intelligence of Low Dimensional Topology 2006*, Eds. J. S. Carter et. al., pp. 101–108, World Scientific Publishing Co., 2007.
- [18] S. Kamada and K. Oshiro, Homology groups of symmetric quandles and cocycle invariants of links and surface-links, *Trans. Amer. Math. Soc.* **362** (2010), no. 10, 5501–5527.
- [19] A. Kawauchi, T. Shibuya and S. Suzuki, Descriptions on surfaces in four-space, I Normal forms, *Math. Sem. Notes Kobe Univ.* **10** (1982), 75–125.
- [20] A. Kawauchi, *A survey of knot theory*, Birkhäuser, 1996.
- [21] C. Kearton and V. Kurlin, All 2-dimensional links in 4-space live inside a universal 3-dimensional polyhedron, *Algebr. Geom. Topol.* **8** (2008), 1223–1247.
- [22] J. Kim, Y. Joung and S. Y. Lee, On the Alexander biquandles of oriented surface-links via marked graph diagrams, *J. Knot Theory Ramifications* **23** (2014), no. 7, 1460007 (26 pages).
- [23] J. Kim, Y. Joung and S. Y. Lee, On generating sets of Yoshikawa moves for marked graph diagrams of surface-links (preprint), 2014, arXiv:1412.5288 [math.GT]
- [24] S. Y. Lee, Invariants of surface links in $\mathbb{R}^4$ via classical link invariants, in *Intelligence of low dimensional topology 2006*, Series on Knots Everything, Vol. **40**, World Scientific Publishing, Hackensack, NJ, 2007, 189–196.

- [25] S. Y. Lee, Invariants of surface links in $\mathbb{R}^4$ via skein relation, *J. Knot Theory Ramifications* **17** (2008), 439–469.
- [26] S. Y. Lee, Towards invariants of surfaces in 4-space via classical link invariants, *Trans. Amer. Math. Soc.* **361** (2009), 237–265.
- [27] S. J. Lomonaco, Jr., The homotopy groups of knots I. How to compute the algebraic 2-type, *Pacific J. Math.* **95** (1981), 349–390.
- [28] S. V. Matveev, Distributive groupoids in knot theory. (Russian) *Mat. Sb. (N.S.)* 119 (161) (1982), no. 1, 78–88; *Math. USSR-Sb.* **47** (1984), 73–83 (in English).
- [29] D. Roseman, Reidemeister-type moves for surfaces in four dimensional space, *Banach Center Publications* **42** (1998) Knot theory, 347–380.
- [30] M. Soma, Surface-links with square-type ch-graphs, Proceedings of the First Joint Japan-Mexico Meeting in Topology (Morelia, 1999), *Topology Appl.* **121** (2002), 231–246.
- [31] F. J. Swenton, On a calculus for 2-knots and surfaces in 4-space, *J. Knot Theory Ramifications* **10** (2001), 1133–1141.
- [32] K. Yoshikawa, An enumeration of surfaces in four-space, *Osaka J. Math.* **31** (1994), 497–522.